			//- 414 Rec'd PC	T/PTO 1-5 NOV 2000 T 8
ORM I	PTO-139	0 (Modified) U.S. DEPARTME	NT OF COMMERCE PATENT AND TRADEMARK OFFICE	ATTORNEY'S DOCKET NUMBER
EV I		RANSMITTAL LETTE	R TO THE UNITED STATES	PK9857
		DESIGNATED/ELEC	TED OFFICE (DO/EO/US)	U.S. APPLICATION NO. (IF KNOWN, SEE 3.7 CER
			ING UNDER 35 U.S.C. 371	To be assign $9/700602$
ITE.	RNAT	IONAL APPLICATION NO.	INTERNATIONAL FILING DATE	PRIORITY DATE CLAIMED
		PCT/JP99/02698	May 24, 1999	May 22, 1998
		NVENTION	Novel Characteristics and Method for Se	narating Substances
		ographic Facking Having I the Same	Novel Characteristics and internod for Se	parating Substances
			010	,
		T(S) FOR DO/EO/US Ikno, et al.		
. • • •	10 04	ikilo, et al.	( NOV 1 5 2000 S)	
nnl	icant l	perewith submits to the United	States Designat Elected Office (DS/EO/US) t	he following items and other information:
••				<u> </u>
1.	$\boxtimes$		of items concerning a filing under 35 U.S.C. 371	
2.			EQUENT submission of items concerning a filing	
3.	X	This is an express request to be examination until the expiration	begin national examination procedures (35 U.S.Co on of the applicable time limit set in 35 U.S.C. 3	C. 371(f)) at any time rather than delay 371(b) and PCT Articles 22 and 39(1).
4.	X			2 19th month from the earliest claimed priority date.
5.	$\boxtimes$	• •	pplication as filed (35 U.S.C. 371 (c) (2))	
٠.		• • •	ith (required only if not transmitted by the Inter	national Bureau)
			by the International Bureau.	610
			e application was filed in the United States Reco	eiving Office (RO/US).
5.	$\boxtimes$		nal Application into English (35 U.S.C. 371(c)(	
7.		A copy of the International Se	- ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' '	MUT 1 3 ZUM 50
8.	$\boxtimes$	• •	the International Application under PCT Article	e 19 (35 U.S.C. 371 (c)(3))
		a.   are transmitted herev	with (required only if not transmitted by the Inte	rnational Bureau).
		b.   have been transmitte	ed by the International Bureau.	
		c.   have not been made;	however, the time limit for making such amend	ments has NOT expired.
		d. 🛮 have not been made	and will not be made.	
9.		A translation of the amendme	nts to the claims under PCT Article 19 (35 U.S.	C. 371(c)(3)).
0.		An oath or declaration of the	inventor(s) (35 U.S.C. 371 (c)(4)).	
1.		A copy of the International Pr	eliminary Examination Report (PCT/IPEA/409)	
2.		A translation of the annexes to (35 U.S.C. 371 (c)(5)).	the International Preliminary Examination Rep	port under PCT Article 36
T4	tome 1		ent(s) or information included:	
3.			tatement under 37 CFR 1.97 and 1.98.	
4.			recording. A separate cover sheet in compliance	with 37 CFR 3.28 and 3.31 is included.
5.	$\boxtimes$	A FIRST preliminary amendr		
6.		A SECOND or SUBSEQUE		
7.		A substitute specification.		
8.		A change of power of attorney	y and/or address letter.	
9.	$\boxtimes$	Certificate of Mailing by Expr		
0.	X	Other items or information:		
		Return Postcard		
		A copy of the International	Application as published by the International	Bureau
		Duplicate copy of this transr	nittal for charging purposes	

529 Rec'd PCT/PTC 15 NOV2000

U.S. APPLICATI	ON NO (IF KNOWN, SEE 37 GER 2	ATTORNEY'S DOCKET NUMB PK9857									
21. The	e following fees are submitted:.	CALCULATION	S PTO USE ONLY								
BASIC NATIO	ONAL FEE ( 37 CFR 1.492 (a) (1) -	* **									
□ Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2) paid to USPTO and International Search Report not prepared by the EPO or JPO											
☑ Internati USPTO											
☐ International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO											
but all c											
☐ International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4)											
0 1 001		ATE BASIC FEE AMO			\$860.00						
months from the	30.00 for furnishing the oath or declar earliest claimed priority date (37 Claimed priority date)	FR 1.492 (e)).			\$0.00						
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		co oo						
Total claims	14 - 20 =	0	x \$18.0 x \$78.0		\$0.00 \$0.00						
Independent cla	ims 3 - 3 = ndent Claims (check if applicable).	<u> </u>	X \$78.0	0	\$0.00						
Multiple Deper	······································	ABOVE CALCULAT		=	\$860.00						
Reduction of 1/2 must also be fil	2 for filing by small entity, if applicated (Note 37 CFR 1.9, 1.27, 1.28) (ch	ble. Verified Small Entity State			\$0.00	:					
a.,		SUB	TOTAL	=	\$860.00						
Processing fee of months from the	of \$130.00 for furnishing the English e earliest claimed priority date (37 Cl	translation later than   20		) +	\$0.00						
		TOTAL NATIONAL	L FEE	=	\$860.00						
Fee for recordin	g the enclosed assignment (37 CFR 1) an appropriate cover sheet (37 CFR	.21(h)). The assignment must b 3.28, 3.31) (check if applicable	e).		\$0.00						
**		TOTAL FEES ENCL	OSED	=	\$860.00						
					Amount to be: refunded	\$					
======================================					charged	\$					
A check	k in the amount of	to cover the above fees is enc	losed.								
1	charge my Deposit Account No.	500-588 in the amount of	\$860.00		to cover the above	ve fees.					
11 449	noute copy of this shoet is shoulded.										
	mmissioner is hereby authorized to cosit Account No. 500-588	harge any fees which may be req A duplicate copy of this sheet is	•	edit an	y overpayment						
NOTE: Where	an appropriate time limit under 37	7 CFR 1.494 or 1.495 has not b	een met, a p	petitio	n to revive (37 CFR	t					
1.137(a) or (b))	must be filed and granted to restor	e the application to pending st	atus.								
	RRESPONDENCE TO:		Lore	//	Monry	erou /					
	armacia Biotech, Inc.		SIGNATI	JRE							
800 Centennia Piscataway, N	ew Jersey 08855		Royal N.	. Ron	ning, Jr.						
(732) 457-8423	•		NAME								
			32,529	ATIO	N NUMBER						
			Novemb	er 15,	2000						
DATE											

09/700602 529 Rec'd PCT/TT 15 NOV 2000 PK-9857

# IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Application of:

Teruo Okano, et al.

Group Art Unit: To be assigned

Serial Number:

To be assigned

Examiner: To be assigned

Filing Date:

15 November 2000

For: Chromatographic Packing Having Novel Characteristics and Method for Separating Substances by Using the Same

## PRELIMINARY AMENDMENT

Honorable Assistant Commissioner of Patents and Trademarks Washington, D.C. 20231

Sir:

Please consider the following amendments and remarks in connection with the prosecution of the captioned application, which claims priority to application PCT/JP99/02698 filed May 24, 1999.

#### IN THE CLAIMS

In Claim 5, line 1, please delete "or 4", without prejudice.

In Claim 7, line 1-2, please delete "claims 1 to 6" and substitute -- Claim 1-- therefor.

In Claim 10, line 1, please delete "or 9", without prejudice.

In Claim 11, line 1-2, please delete "claims 8 to 10" without prejudice, and substitute --Claim 8--therefor.

In Claim 14, line 1, please delete "or 13" without prejudice

#### **REMARKS**

Claims 1-14 are pending in the captioned application.

Applicants have amended Claims 5, 7, 10, 11, and 14 to delete multiple dependencies.

Applicants respectfully submit that the amendments are fairly based on the specification, and respectfully request their entry.

Applicants believe that the claims, as amended, are in allowable form, and earnestly solicit the allowance of claims 1-14.

Respectfully submitted,

Royal N. Ronning, Jr. 32,529

Attorney for Applicants

Amersham Pharmacia Biotech Inc. 800 Centennial Avenue P. O. Box 1327 Piscataway, NJ 08855-1327

Tel: (732) 457-8423

RR/mp

H:\DEPT\Law\_Dept\IP Dept\US Patent Forms PDF & Word\Word Forms\Preliminary Amendment-PK9857 doc

PK9857/PCT/JP99/02698/YCT-424 PCT Application (Translation)

SPECIFICATION

# CHROMATOGRAPHIC PACKING HAVING NOVEL CHARACTERISTICS AND METHOD FOR SEPARATING SUBSTANCES BY USING THE SAME

5

10

#### Technical Field

This invention relates to a packing which contains a charged (co)polymer and makes it possible to change the effective charge density or hydrophilic/hydrophobic balance on the surface of a stationary phase in an aqueous system by an external signal (for example, temperature), and a novel separation method by which substances such as metal elements, drugs or biological components are chromatographically separated by using the packing.

15

20

25

#### Background Art

There a great variety of liquid chromatography techniques depending on the combination of stationary phase with mobile phase and the interaction systems employed for the separation. Liquid chromatography is a highly important technique for separating metal elements, isolating and purifying drugs and separating peptides, proteins, nucleic acids, etc. in the field of biochemistry. In recent years, moreover, attempts have been made to apply recombinant proteins, etc. produced by bioengineering procedures, which have made remarkable advances, to medicines. Under these circumstances, there is a growing requirement for efficient separation methods for separating and purifying these products. Chromatographic techniques commonly employed at present involve ion-exchange chromatography,

10

15

20

25

reversed phase chromatography, etc.

In ion-exchange chromatography, separation is carried out by using, as a stationary phase, an electrolyte on the surface of an insoluble carrier and irreversibly adsorbing counter ions contained in the mobile phase. As the carrier, silica, cellulose, dextran, styrene/divinylbenzene copolymer, etc. are widely employed. Carriers having ion-exchange groups (for example, sulfonate, quaternary ammonium) introduced thereinto are commercially available as ion exchangers. Solute dissociate into cations, anions and amphoteric ions depending on the hydrogen ion concentration in the solution. When this solution is passed through an ion-exchange column, each ion binds to the oppositely charged exchange group on the carrier surface competitively with solvent ions, thus causing distribution between the solution and the ion exchanger surface at a certain ratio. The migration rates through the column vary depending on the bond strength and separation is completed by utilizing this difference in the migration rate. The distribution can be modified by some methods. For example, it can be changed by controlling the concentration of the competitive ion species in the mobile phase. Alternatively, the extent of ionization of the ion-exchange group on the carrier surface may be varied by changing the hydrogen ion concentration in the solution. That is to say, it has been a practice in ion-exchange chromatography to separate solutes from each other by controlling the ionic strength or the hydrogen ion concentration in the mobile phase to thereby change the elution order of the solutes.

Reversed phase chromatography involves the use of a combination of a hydrophobic stationary phase and a polar mobile

5

10

15

20

25

Ŷŗ

phase. Solutes are distributed between the mobile phase and the stationary phase depending on the degree of hydrophobicity. In this case, solutes are eluted also by changing the degree of hydrophobicity of the solvent in the mobile phase to thereby change the distribution between the mobile phase and the stationary phase. Since an organic solvent is employed as the solvent in the mobile phase, it is feared that the activities of the biological components to be separated might be caused to deteriorated thereby.

In short, solutes are eluted and separated from each other fundamentally by varying the solvent in the mobile phase both in ion-exchange chromatography and reversed phase chromatography. Accordingly, there is a risk that the activity of the target sample might be damaged by an acid or organic solvent employed in the elution.

When it is intended to separate substances from each other by two or more chromatographies, each chromatography should be independently carried out, since chromatographic mode varies from carrier to carrier. If it is possible to perform ion-exchange chromatography and reversed phase chromatography by using a single carrier and a single physical stimulus, separation could be completed at an elevated efficiency within a shorter period of time. Moreover, substances which cannot be separated from each other by the conventional techniques can be separated thereby.

There are a great variety of biological components including charged ones and uncharged ones. In general, a compound capable of being ionized is retained, in an unionized state, in a hydrophobic packing owing to hydrophobic interaction.

r i

5

10

15

20

25

When ionized, however, the hydrophobic interaction with the hydrophobic packing is weakened. Ion-dissociatable compounds differing in the dissociation constant can be easily separated from each other owing to the ion-ion interaction with the use of an ion exchanger.

It is generally known that weakly acidic ion exchange resins and weakly basic ion exchange resins are suitable respectively for separating basic proteins and acidic proteins. It is thus expected that, by introducing ion-exchange substituents, ion-exchange chromatography based on ion-ion interactions becomes usable in separating various substances, which are similar to each other in hydrophobicity or molecular weight and thus cannot be separated exclusively by hydrophobic interactions, and biological molecules such as proteins and nucleic acid oligomers.

However, there has been known hitherto neither any carrier which is usable both in ion-exchange chromatography and reversed phase chromatography when employed alone under one physical stimulus nor one usable in efficiently separating various substances, which are similar to each other in hydrophobicity or molecular weight and thus cannot be separated exclusively by hydrophobic interactions, and biological molecules such as proteins and nucleic acid oligomers.

#### Disclosure of Invention

To solve the above-mentioned problems, the present inventors have conducted studies and developments from various viewpoints. As a result, they have successfully prepared a novel packing having ion-exchange function by copolymerizing

5

10

15

20

25

poly(N-isopropylacrylamide)(PIPAAm) with positively charged dimethylaminopropylacrylamide (DMAPAAm) and found that this packing is usable both in reversed phase chromatography and ion-exchange chromatography, when temperature is properly controlled. They have furthermore found that use of the charged copolymer makes it possible to control the LCST of the polymer by regulating pH value. The present invention has been completed based on these findings.

The present invention relates to a method for separating substances characterized by chromatographically separating said substances with the use of a packing which contains a charged (co)polymer and makes it possible to change the effective charge density on the surface of a stationary phase by an external stimulus while fixing a mobile phase to an aqueous system.

The present invention further relates to a method for separating substances characterized by retaining the substances in a stationary phase made of a chromatographic packing chemically modified with a polyalkylacrylamide copolymer having amino, carboxyl, hydroxyl groups, etc., then changing the hydrophilic/hydrophobic balance on the surface of the stationary phase by the temperature gradient method wherein the external temperature is changed stepwise, and passing the substances through a single mobile phase to thereby separate the same.

The present invention furthermore relates to a chromatographic packing which contains a charged (co)polymer and makes it possible to change the effective charge density on the surface of a stationary phase by a physical stimulus.

In the chromatographic packing of the present invention,

10

15

20

25

the charged state of ion-exchange groups on the surface of a carrier can be reversibly controlled by changing the surface structure of the stationary phase by an external physical stimulus such as a change in temperature. Namely, the present invention provides a stationary phase which makes it possible to perform two chromatographic modes, i.e., ion-exchange chromatography and reversed phase chromatography, at the same time with the use of a mobile phase which is a single aqueous solvent (aqueous mobile phase). Moreover, the present invention provides a carrier capable of arbitrarily controlling the charge of ion-exchange groups on the surface of the carrier (in the case of ion-exchange chromatography) or the hydrophilic/hydrophobic balance (in the case of the reversed phase chromatography). The term "aqueous solvent" as used herein means water alone or aqueous solutions containing inorganic salts but free from any organic solvent.

The present invention provides a carrier for separation and purification characterized in that separation is performed by controlling the charge of ion-exchange groups on the surface of the stationary phase by regulating the physical properties or structure around the ion exchange groups on the carrier surface by a physical stimulus, while fixing the mobile phase to an aqueous system. According to the present invention, when the external temperature is lower than the critical temperature, the ion-exchange groups appear on the surface of the carrier. Then the biological components to be separated undergo interaction with the ion-exchange groups followed by separation by the ion-exchange chromatography mode. When the external temperature is higher than the critical temperature, on the other

10

15

20

25

hand, the surface charge is weakened and the carrier becomes more hydrophobic. Then, the biological components can be separated by the reversed phase chromatography mode. That is to say, the hydrophilic/hydrophobic balance on the surface of the carrier can be reversibly and arbitrarily changed by controlling the external temperature.

#### Brief Description of Drawings

Fig. 1 provides graphs showing the relationship between temperature and retention time in the separation of aspirin, salicylic acid, methyl salicylate and benzoic acid with the use of two packings described in Example 2.

Fig. 2 provides graphs showing the relationship between temperature and retention time in the separation of aspirin, salicylic acid, methyl salicylate and benzoic acid while changing pH value in Example 3.

Fig. 3 provides graphs showing the relationship between temperature and retention time in the separation of aspirin, salicylic acid, methyl salicylate and benzoic acid while changing ionic strength in Example 4.

Fig. 4 provides graphs showing the relationship between temperature and retention time in the separation of aspirin, salicylic acid, methyl salicylate and benzoic acid while changing the polymerization ratio of IPAAm to DMAPAAm in Example 5.

#### Best Mode for Carrying Out the Invention

The external physical signal to be used in the method of the present invention is exemplified by a change in temperature.

10

15

20

25

To alter the physical properties or structure around ionexchange groups on the surface of the packing by changing
temperature, for example, a temperature-responsive polymer may
be introduced into the surface of the carrier. Examples of
packings of this type include chromatographic packings
chemically modified on the surface of the carrier with
alkylacrylamide polymers or copolymers having amino, carboxyl,
hydroxyl groups, etc. in the side chains or at the ends.
Chemically modified packings are exemplified by silica carriers
modified with the above-mentioned alkylacrylamide polymers or
copolymers. To introduce ion-exchange groups, carriers may be
chemically modified by copolymers of the above-mentioned
alkylacrylamides with comonomers having amino or carboxy
groups.

Examples of the constitutional units of amino-containing polymers include dialkylaminoalkyl(meth)acrylamide, dialkylaminoalkyl (meth)acrylate, aminoalkyl (meth)acrylate, aminostyrene, aminoalkylstyrene, aminoalkyl(meth)acrylamide, alkyloxyalkyltrimethylammonium salts and (meth)acrylamido-alkyltrimethylammonium salts. Examples of the constitutional units of carboxyl-containing polymers include acrylic acid and methacrylic acid, while examples of the constitutional units of the sulfonate-containing polymers include (meth)acrylamido-alkylsulfonic acid.

It is preferable that the polyalkylacrylamide to be used in the present invention is selected from among poly(N-isopropylacrylamide), polydiethylacrylamide, poly(N-propylacrylamide) and polyacryloylpyrrolidine and copolymers of the constitutional units of these polymers with alkyl

(meth)acrylate, as shown by the following formulae.

# [Chemical formula 1]

## Polyalkylacrylamide

5

 $CH_2-CH$  C=0  $R_1$   $R_2$ 

10

15

 $R_1$   $R_2$  Abbreviation

poly(N,N'-diethylacrylamide)  $-C_2H_5$   $-C_2H_5$  poly(DEAAm)

20

poly(N-propylacrylamide) -H -C<sub>3</sub>H<sub>7</sub> poly(PAAm)

25

10

15

#### [Chemical formula 2]

#### Copolymer

$$\begin{array}{c|c}
\hline
 & \text{CH}_2\text{-CH} \\
 & \text{C} \\
 & \text{C} \\
 & \text{C} \\
 & \text{N}
\end{array}$$

A: content: 50 - 60 %

Alkyl acrylate (t = 1 - 20)

$$\begin{array}{c}
B \\
-CB_2 - C - \\
COOC_tB_{2t+1}
\end{array}$$
Alkyl methacrylate (t = 1 - 20)

$$\begin{array}{c}
CB_3 \\
-CB_2 - C - \\
COOC_tB_{2t+1}
\end{array}$$

Since poly(N-isopropylacrylamide) has a lower limit of critical temperature of 32 °C, a carrier chemically modified therewith undergoes a large change in the hydrophilic/ hydrophobic surface properties at this critical temperature. When the surface of a chromatographic packing is grafted or coated with this polymer, the power of retaining a sample varies depending on temperature. Thus, the retention behavior can be regulated by controlling temperature without changing the composition of the eluate. A lower limit of critical temperature of 32 °C or above can be achieved by copolymerizing

the N-isopropylacrylamide with comonomers which are more

10

15

20

25

hydrophilic than isopropylacrylamide, for example, acrylamide, methacrylic acid, acrylic acid, dimethylacrylamide and vinyl pyrrolidone. On the other hand, a lower limit of critical temperature lower than 32 °C can be achieved by copolymerizing the N-isopropylacrylamide with hydrophobic comonomers, for example, styrene, alkyl methacrylate and alkyl acrylate.

The lower limit of critical temperature of polydiethylacrylamide is about 30 to 32 °C. At this temperature, this polymer undergoes a change in the surface hydrophilic/hydrophobic nature. Similar to the abovementioned case of poly(N-isopropylacrylamide), the power of retaining a sample can be thus regulated by controlling temperature. The novel chromatographic carrier to be used in the present invention is prepared by chemically modifying or coating the carrier with a polymer. The chemical modification can be carried out by two methods, i.e., surface grafting and radical polymerization. In the case of coating, on the other hand, the polymer is insolubilized within the application temperature range and then the insolubilized product is employed in coating.

As described above, surface grafting and radical polymerization can be employed as the chemical modification means by which a temperature-responsive polymer is introduced into a carrier. In the surface grafting method, a temperature-responsive polymer of a definite size is first synthesized and then grafted to the carrier. In the radical polymerization method, in contrast thereto, monomer(s) are polymerized on the surface of the carrier to give a polymer. Compared with the surface grafting method, the radical

10

15

20

polymerization method makes it possible to introduce the temperature-responsive polymer into the surface of the carrier at a high density. Thus, the hydrophobicity of the surface of the carrier can be elevated and the retention time can be easily controlled. In this case, moreover, non-specific adsorption on the carrier surface due to the interaction with silica gel can be easily suppressed.

Substances which can be separated by the method of the present invention include metal element (Cu<sup>2+</sup>, Mn<sup>2+</sup>, etc.), drugs (steroids, antipyretic analgesic agents, etc.) and biological components (peptides, proteins, nucleic acids, etc.). The method of the present invention is particularly useful in separating various biological components which cannot be separated by using either ion-exchange chromatography or reversed phase chromatography alone.

#### <u>Examples</u>

To further illustrate the present invention in greater detail, and not by way of limitation, the following Examples will be given.

#### [Example 1]

- 1. Synthesis of polymer
- 1-1) Poly(IPAAm-DMAPAAm) (DMAPAAm:N,N-dimethylaminopropyl-acrylamide)
- 25 1-1-a) Preparation of IPAAm copolymer having carboxyl end
  An IPAAm copolymer having a carboxyl end was synthesized
  in such a manner as to give a molecular weight of 4,000 as a
  standard. The molecular weight of the polymer can be designed
  by controlling the amount of 3-mercaptopropionic acid (MPA)

15

20

25

employed as a chain transfer agent. To prepare a copolymer having a molecular weight of 4,000, the amount of MPA was regulated so as to give a molar ratio MPA/(IPAAm + DMAPPAm) of 0.028.

5 Purified monomer IPAAM: 25.0 g.

Cationic monomer (5 % by mol of DMAPPAm based on IPAAm):

Radical polymerization initiator [2,2'-azobis(isobutyro-nitrile) (AIBN): 0.145 g.

Chain transfer agent (3-mercaptopropionic acid): 0.691 g.

DMF (N,N-dimethylformamide): 50 ml.

The above components were fed into a polymerization tube and fixed with a rubber ring provided with a three-way stopcock. The polymerization tube was introduced into liquid nitrogen, while closing the cock, and completely frozen. Next, the cock was opened and the contents of the tube were degassed by using a vacuum pump. After closing the cock again, the polymerization tube was introduced into methanol and the sample in the tube was completely dissolved. This procedure was repeated thrice (freezing/thawing degassing method). Then the polymerization tube containing the completely degassed sample under reduced pressure was introduced into a thermostat under shaking at 70 °C and radical polymerization was performed for 2 hours to thereby synthesize a copolymer having a carboxyl group at one end. After the completion of the reaction, the reaction mixture was cooled to room temperature by allowing to stand. Then the solvent (DMF) was concentrated by distilling at 40 °C under reduced pressure and the residue was dropped into ice-cooled diethyl ether to thereby give a polymer. The polymer thus obtained was taken up

10

15

20

25

by filtration and dried at ordinary temperature under reduced pressure overnight. The dried product was dissolved in acetone and purified again with diethyl ether. The polymer thus obtained was taken up again by filtration and dried at ordinary temperature under reduced pressure overnight. The obtained polymer was then dissolved in purified water to give a 5 % (w/v) solution. The resultant solution was transferred onto a dialysis membrane of a fractional molecular weight of 500 and dialyzed for 3 days. Thus a highly pure copolymer having a uniform molecular weight could be obtained.

- 1-1-b) Introduction of IPAAm copolymer into carrier
- (a) Active esterification (succinylation) method

To succinylate the copolymer synthesized above, the molar ratio of the copolymer: N,N'-dicyclohexylcarbodiimide (DCC): N-hydroxysuccinimide was adjusted to 1 : 2.5 : 2.

The copolymer was fed into a round-bottomed flask and dissolved in a half amount (25 to 30 mL) of ethyl acetate. Next, N-hydroxysuccinimide and DCC were added thereto followed by dissolution in the residual ethyl acetate. The obtained mixture was immersed in ice-water at 4 °C and stirred with a stirrer for 2 hours. Subsequently, it was introduced into a thermostat at 25 °C and stirred therein overnight. The solution was filtered and thus dicyclohexyl urea formed as a by-product was removed therefrom. After concentrating under reduced pressure, the residue was purified with diethyl ether. The product thus obtained was taken up by filtration and dried under reduced pressure. The succinylated copolymer thus obtained was stored in a freezer.

1-1-c) Introduction into carrier (silica gel)

10

The succinylated copolymer was reacted in three portions with aminopropyl silica gel with the use of 1,4-dioxane as a solvent. The reaction was carried out at room temperature (25°C). First, the succinylated polymer (1.0 g) was dissolved in 1,4-dioxane (50 mL) and reacted with aminopropyl silica gel (3 g) in a thermostat under shaking overnight. Subsequently, the liquid reaction mixture was filtered and the precipitate thus obtained and fresh copolymer (1.0 g) were dissolved in 1,4-dioxane (50 mL) again and reacted overnight. After repeating this procedure once again, the product finally taken up by filtration was sufficiently washed with methanol (500 mL) and distilled water (2 mL), dried under reduced pressure and then stored in a desiccator as a packing.

15 [Example 2]

- 1-2) Preparation of PIPAAm hydrogel surface
- 1-2-a) Formation of gel layer on aminopropyl silica gel surface

To introduce a polymerization initiator into aminopropyl silica gel, the following compounds were used.

Aminopropyl silica gel: 5 g.

V-501 : 3.5 g (12.5 mmol).

EEDQ : 6.18 g (25.0 mmol).

25 DMF : 50 ml.

Use was made of V-501 [4,4'-azobis(4-cyanovaleric acid) (molecular weight: 280.28)] as a polymerization initiator and EEDQ [N-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline,

20

25

5

molecular weight: 247.30] as a condensing agent each in the amount as specified above. These compounds were reacted with aminopropyl silica gel in DMF. After bubbling  $N_2$  gas thereinto in the dark for 30 minutes, the reaction vessel was completely charged with  $N_2$  and reaction was carried out by using an  $N_2$  balloon at room temperature for 6 hours. After the completion of the reaction, the mixture was filtered and washed with DMF. Thus, the polymerization initiator had been introduced into the surface of the aminopropyl silica gel.

10 1-2-b) Formation of surface gel layer

Silica gel having V-501 bonded thereto prepared

in above 1-2-a): 4 g.

IPAAm : 10 g.

BIS : 0.27 g.

EtOH : 200 ml.

DMAPAAm : such an amount as to give a molar

ratio to IPAAm of 8:2 or 9:1.

Silica gel, IPAAm, DMAPAA and BIS [N,N'-methylene-bis (acrylamide), molecular weight: 154.17] were dissolved in ethanol. After bubbling N<sub>2</sub> gas thereinto in the dark for 1 hour, the reaction vessel was completely charged with N<sub>2</sub> and reaction was carried out in an oil bath at 70 °C by using an N<sub>2</sub> balloon for 5 hours, thus forming a gel layer on the surface of PIPAAm. After the completion of the reaction, the mixture was filtered and washed with methanol and water. The obtained product was dried under reduced pressure and stored in a desiccator as a packing. It was packed into a stainless column and employed in

analysis.

#### [Example 3]

5

10

20

25

Aspirin, salicylic acid, methyl salicylate and benzoic acid were separated under the following conditions by using columns packed with a positively charged gal (IPAAm: DMAPAAm = 8:2) and IPAAm hydrogel.

#### Separation conditions

Column: (1) packed with poly(IPAAm) hydrogel-modified silica;

(2) packed with poly(IPAAm-co-DMAPAAm) (8 : 2)
hydrogel-modified silica.

Buffer : Na<sub>2</sub>CO<sub>3</sub>/NaHCO<sub>3</sub>.

pH = 9.0.

15 Ionic strength = 0.1 M.

Fig. 1 shows the results. Aspirin could not be separated from benzoic acid by using the column packed with the IPAAm hydrogel. In contrast, these compounds could be separated from each other by using the column packed with the positively charged gel (IPAAm: DMAPPAAm = 8:2). At 10 °C, in particular, all of the four compounds including charged and uncharged ones could be separated from each other within a short period of time of about 20 minutes. The order of separation depended on the hydrophobicity degrees of these compounds. In the cases of salicylic acid and benzoic acid, the retention times were shortened as temperature was elevated. This is seemingly because, when temperature was elevated, the structure and physical properties of the temperature-responsive polymer were changed and the charge on the carrier surface was thus lowered

10

15

20

25

so as to reduce the interactions between the surface and the solutes. On the contrary, methyl salicylate (i.e., an uncharged compound) showed an prolonged retention time as temperature was elevated. This is seemingly because the temperature-responsive polymer became hydrophobic due to increase in temperature.

## [Example 4] Effects of pH change

Aspirin, salicylic acid, methyl salicylate and benzoic acid were separated by the same method as the one of Example 3 but using the column packed with poly(IPAAm-co-dMAPAAm) (8:2) hydrogel-modified silica of Example 3 and NaHPO<sub>4</sub>/H<sub>3</sub>C<sub>6</sub>H<sub>5</sub>O<sub>7</sub> [citric acid H<sub>2</sub>O (monohydrate)] as a buffer at pH 7.0. Fig. 2 shows the results.

Fig. 2 indicates that the retention times of all of the substances were prolonged at pH 7.0, compared with at pH 9.0. This is seemingly because anionic compounds would undergo stronger interactions with the positively charged carrier surface at pH 7.0. These results suggest that the retention times of substances to be separated can be controlled by regulating pH value.

#### [Example 5] Effects of ionic strength

Aspirin, salicylic acid, methyl salicylate and benzoic acid were separated by using a column packed with the poly(IPAAm-co-DMAPAAm) (8:2) hydrogel-modified silica of Example 3 under the following separation conditions.

Separation conditions

Buffer :  $NaHPO_4/H_3C_6H_5O_7$ . pH = 7.0.

10

15

20

Ionic strength = 1.0 M and 0.1 M.

Fig. 3 shows the results. As the ionic strength was elevated (0.1 M  $\rightarrow$  1.0 M), the retention times of all of the compounds but the uncharged methyl salicylate were shortened, while the retention time of methyl salicylate was prolonged. This is seemingly because, when the ionic strength was elevated, the protonation of amino groups on the surface of the carrier was suppressed and the positive charge was lowered, which weakened the interactions of the carrier surface with the anionic compounds. In the case of methyl salicylate, the hydrophobicity was elevated with an increase in the ionic strength and, in its turn, the hydrophobic interaction was seemingly strengthened.

[Example 6] Effect of polymerization ratio of IPAAm to DMAPAAm Aspirin, salicylic acid, methyl salicylate and benzoic acid were separated under the following conditions.

Separation conditions

- Column: (1) packed with poly(IPAAm-co-DMAPAAm) (9:1) hydrogel-modified silica;
  - (2) packed with poly(IPAAm-co-DMAPAAm) (8 : 2)
    hydrogel-modified silica.

Buffer :  $NaHPO_4/H_3C_6H_5O_7$ .

pH = 7.0.

25 Ionic strength = 0.1 M.

Fig. 4 shows the results. The retention times were prolonged with an increase in the ratio of the positively charged polymer, which indicates that retention time can be controlled by regulating the polymerization ratio.

10

15

20

#### Industrial Applicability

The present invention has the following advantages.

- 1) The charge of an ion exchanger exposed on the surface of the carrier can be arbitrarily controlled by regulating temperature. Thus separation can be performed in a single aqueous mobile phase without changing the solvent in the mobile phase.
- 2) Due to differences in hydrophobicity and ionic properties, separation can be carried out by a single operation. Compared with the conventional methods wherein two separating operations are needed, therefore, the method of the present invention is a highly efficient one and gives an elevated yield.
- 3) The method of the present invention makes it possible to separate biological components which cannot be separated by either ion-exchange chromatography or reversed phase chromatography employed alone.
- 4) Since neither any acid nor organic solvent is used in the method of the present invention, biological components can be separated without deteriorating their activities.
- 5) Compared with the conventional ion exchangers, the packing of the present invention can be quickly regenerated.

25

#### Claims

- A method for separating substances characterized by chromatographically separating said substances with the use of a packing which contains a charged (co)polymer and makes it possible to change the effective charge density on the surface of a stationary phase by a physical stimulus while fixing a mobile phase to an aqueous system.
- The separation method as claimed in Claim 1, wherein said
   physical stimulus is a change in temperature.
  - 3. The separation method as claimed in Claim 2, wherein said packing is a chromatographic packing chemically modified on the surface of a carrier with a temperature-responsive polymer.
  - 4. The separation method as claimed in Claim 3, wherein said packing is a chromatographic packing chemically modified with a temperature-responsive polymer by using the radical polymerization method.
- 5. The separation method as claimed in Claim 3 or 4, wherein said temperature-responsive polymer, with which the surface of the carrier is chemically modified, is a polyalkylacrylamide polymer or copolymer having amino, carboxyl, hydroxyl groups, etc. in the side chains or at the ends.
  - 6. The separation method as claimed in Claim 5, wherein said polyalkylacrylamide is one selected from among poly(N-isopropylacrylamide), poly(N-propylacrylamide),
  - polydiethylacrylamide and polyacryloylpyrrolidine.
    - 7. The separation method as claimed in any of Claims 1 to 6, wherein said substances are those selected from among metal elements, drugs and biological components.

: 14 d

- 8. A method for separating substances characterized by retaining the substances in a stationary phase made of a chromatographic packing chemically modified with a polyalkylacrylamide copolymer having amino, carboxyl, hydroxyl
- 5 groups, etc., then changing the hydrophilic/
  hydrophobic balance on the surface of the stationary phase by
  the temperature gradient method wherein the external
  temperature is changed stepwise, and passing the substances
  through a single mobile phase to thereby separate the same.
- The separation method as claimed in Claim 8, wherein said mobile phase is an aqueous solvent.
  - 10. The separation method as claimed in Claim 8 or 9, wherein said polyalkylacrylamide is one selected from among poly(N-isopropylacrylamide), poly(N-propylacrylamide),
- 15 polydiethylacrylamide and polyacryloylpyrrolidine.
  - 11. The separation method as claimed in any of Claims 8 to 10, wherein said substances are those selected from among metal elements, drugs and biological components.
- 12. A chromatographic packing which contains a charged
  20 (co)polymer and makes it possible to change the effective charge density on the surface of a stationary phase by a physical stimulus.
  - 13. The chromatographic packing as claimed in Claim 12, wherein said (co)polymer is a polyalkylacrylamide copolymer having amino, carboxyl, hydroxyl groups, etc.
  - 14. The chromatographic packing as claimed in Claim 12 or 13, wherein said polyalkylacrylamide is one selected from among poly(N-isopropylacrylamide), poly(N-propylacrylamide), polydiethylacrylamide and polyacryloyl-pyrrolidine.

#### Abstract

[Abstract]

[Object] To provide chromatographic packings whereby

- biological components, etc. which cannot be separated by either ion-exchange chromatography or reversed phase chromatography employed alone can be efficiently separated without deteriorating their activities.
- [Means for solution] Use is made of a packing which contains

  10 a charged copolymer and makes it possible to change the effective
  charge density on the surface of a stationary phase by a physical
  stimulus while fixing a mobile phase to an aqueous system.
  [Selected figure] None.

Fig. 1





Fig. 2







Fig. 3








Fig. 4





PTO/SB/01 (12-97)

Approved for use through 9/30/00. OMB 0651-0032

Patent and Trademark Office, U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

#### PK-9857 Attorney Docket Number **DECLARATION FOR UTILITY OR** Okano First Named Inventor **DESIGN** PATENT APPLICATION **COMPLETE IF KNOWN** 09 /700,602 (37 CFR 1.63) Application Number 15-Nov-2000 Filing Date ☐ Declaration ☑ Declaration OR To be assigned Submitted Submitted after Initial Group Art Unit Filing (surcharge (37 CFR 1.16 (e)) with Initial To be assigned Filing **Examiner Name** required)

	As a below named inventor, I hereby declare that:										
My residence, post office address, and citizenship are as stated below next to my name.											
I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:											
Chromatographic Packing Having Novel Charactersitics and Method for Separating Substances by Using the Same											
the specification of which (Title of the Invention)  Is attached hereto  OR											
	DAYYY) 11/15/200	0 as Unite	d States Applica	tion Number or P	CT International						
Application Number 0	9/700,602 and w	as amended on (MM/DD/Y	YYY)		(if applicable)						
I hereby state that I have re amended by any amendme	eviewed and understand the ent specifically referred to abo	contents of the above ident	lified specificatio	n, including the c	laıms, as						
	disclose information which is		defined in 37 CF	R 1.56							
I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed											
certificate, or 365(a) of any America, listed below and ha	PCT international applications are also identified below, by	on which designated at lea	st one country	other than the U	nited States of						
certificate, or 365(a) of any America, listed below and ha	PCT international applications are also identified below, by	on which designated at lea	st one country	other than the U r patent or inven ority is claimed Certified Co	nited States of tor's certificate, py Attached?						
certificate, or 365(a) of any America, listed below and ha or of any PCT international a	PCT international applications are also identified below, by application having a filing date.	on which designated at lea checking the box, any forei e before that of the applicat	st one country gn application fo tion on which pre	other than the U r patent or inven ority is claimed	nited States of tor's certificate,						
certificate, or 365(a) of any America, listed below and he or of any PCT international at Prior Foreign Application Number(s)  140722/1998	PCT international application was also identified below, by application having a filing date.  Country  JP  ation numbers are listed on a	on which designated at lea checking the box, any foreign before that of the applicate (MM/DD/YYYY)  05/22/1998	st one country gn application for tion on which price Priority Not Claimed	other than the U r patent or inven porty is claimed  Certified Co YES	py Attached? NO						
certificate, or 365(a) of any America, listed below and he or of any PCT international at Prior Foreign Application Number(s)  140722/1998	PCT international application was also identified below, by application having a filing date.  Country  JP  ation numbers are listed on a under 35 U.S.C. 119(e) of an	on which designated at lea checking the box, any foreign before that of the applicate (MM/DD/YYYY)  05/22/1998	st one country gn application for tion on which price Priority Not Claimed	other than the U r patent or inven porty is claimed  Certified Co YES	py Attached? NO						

[Page 1 of 2]

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231

Q
7,7
j
L
H
-

Please type a plus sign (+) inside this box → +	PTO/SB/01 (12-97) Approved for use through 9/30/00. OMB 0651-0032
Under the Paperwork Poduction Act of 1005, no name	Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contain a valid OMB control number.

# **DECLARATION** — Utility or Design Patent Application

I hereby claim the benefit under 35 U.S.C. 120 of any United States application(s), or 365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. 112, I acknowledge the duty to disclose information which is material to patentiability as defined in 37 CFR 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. Parent Application or PCT Parent

Number

Parent Filing Date
(MM/DD/YYYY)

(if applicable)

U.S. Parent Application or PCT Parent Number					Parent Filing Date Parent Patent Nun (MM/DD/YYYY) (if applicable)				
PCT/J	IP99/02698			05/24/1999					
Additional U.S. or	PCT international applica	tion numbers ar	re listed on	a supplemer	tal priority data	sheet PTO/SE	3/02B attached	hereto.	
As a named inventor. I h	nereby appoint the following onnected therewith:	ractitioner	e) to proceed	te this application	on and to trans				
		OR Registered pra	ctitioner(s)	name/registr	ation number lis	ted below	Number Ba	40	
Nam	1e		tration nber		Nam	ie	Regi	istration Imber	
	***************************************	1100	, inc.				PATENT TRADER	ARK OFFICE	
,									
Additional registere	d practitioner(s) named o	n supplemental	Registere	d Practitioner	Information sha	et PTO/SR/03	C attached hor	eto	
Direct all correspond		er Number							
		Code Label	2284	40	OR	Corres	oondence ado	tress below	
Name									
Address									
Address									
City				State		ZIP			
Country		Telephor				Fax			
	Il statements made here d further that these state mprisonment, or both, un it issued thereon.								
Name of Sole or I	First Inventor:			☐ A petit	ion has been	filed for this	unsigned inv	entor	
Given Nar	me (first and middle [if	any])			Family	Name or Si	urname		
Teruo				Okano					
Inventor's Signature		(		7			Date	Dec. 27	
Residence: City Chiba		State		country Japan				2 <i>000</i> JP	
Post Office Address	6-12-12, K		ai, Ich			JPX	Citizenship		
Post Office Address Chiba 272-0827									
City	State		ZIP			Country			
Additional invento	rs are being named o	n the 2_su	pplement	al Additiona	I Inventor(s) s	heet(s) PTO	/SB/02A atta	ched hereto	

1-00

# **DECLARATION**

# ADDITIONAL INVENTOR(S) Supplemental Sheet Page 1 of 2

Name of Additional Joint Inventor, if any:							d for this	for this unsigned inventor			
Given Nar	ne (first and middle [if any	])		Family Name or Sumame							
<u>Akihik</u> o			,	<u>Kikuch</u> i							
Inventor's Signature	akis	rik	0	2	Jen	eh_	_	Date		Рес 28 2000	
Residence: City	Tokyo state country Japan citizenship JP							P			
Post Office Address	3-26-1-401, Hyakunin-cho, Shinjuku-ku										
Post Office Address	Tokyo 169-0	073	Jap	an							
City		State			ZIP		Country				
Name of Addition	nal Joint Inventor, if ar	ıy:			A petitio	n has been file	d for this	s unsigr	ned inv	entor	
Given Nar	ne (first and middle [if any	1)				Family Nar	ne or S	umame			
Yasuhisa	a .				Sak	<u>urai</u>					
Inventor's Signature	Jamhi Tokyo	sa		80	hur	<u>ر</u> ٔ		Da	te	Dec. 28 2000	
Residence: City	Tokyo_	State			Country Japan Citizenship J				JP		
Post Office Address	3-17-6, Eifuk	u, Sı	ıgina	ami-	-ku	J	pX				
Post Office Address	Tokyo 168-	0064	Ja	par	)		,				
City		State			ZIP		Count	ntry			
Name of Addition	nal Joint Inventor, if ar	ıy:			A petitio	n has been file	d for this	s unsigi	ned inv	entor	
Given Nar	me (first and middle [if any	])		Family Name or Sumame							
<u>Hideko</u>				Kanazawa							
Inventor's Signature	Hideko Kanazawa				wa Date 2000					Dec 28 2000	
Residence: City	Kanagawa	State			Country Japan JOX Cilizenship JP					JP	
Post Office Address	2896-1-303, K	amits	urum	na, S	Sagar	nihara-sh	ni '				
Post Office Address	Kanagawa 2	28-08	02	Jap	an						
City		State			ZIP		Co	ountry			

Burden Hour Statement. This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time, you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO Assistant Commissioner for Patents, Washington, DC 20231.

Dinama Arma a ultra arma (1) maida Abra barra S		
Please type a plus sign (+) inside this box →	7	

PTO/SB/02A (3-97)
sign (+) inside this box -> + Approved for use through 9/30/98. OMB 0651-0032
Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

# **DECLARATION**

# ADDITIONAL INVENTOR(S) Supplemental Sheet Page 2\_ of 2\_

Name of Addition	al Joint Inventor, if any		A petition has been filed for this unsigned inventor							
Given Nar	ne (first and middle [if any])			Family Name or Sumame						
Yoshikazu				Matsushima						
Inventor's Signature	Joshika	0	Matsushime Date 2000							
Residence: City	<u>Kanagawa</u>			Country	Japan	,	Citizensl	<sub>nip</sub> J	Р	
Post Office Address 2-3-21, Ikeda, Kawasaki-ku, Kawasaki-shi										
Post Office Address	Kanagawa 2	10-0	022	Ja	pan					
City		State			ZIP		Country			
Name of Addition	nal Joint Inventor, if any	<b>/</b> :			A petitio	n has been filed	for this	unsign	ed inv	entor
Given Nar	ne (first and middle [if any])					Family Nan	ne or Su	mame		
Inventor's Signature				Date						
Residence: City		State		Country Citizenship						
Post Office Address										
Post Office Address										
City		State			ZIP		Count	ry		
Name of Additior	nal Joint Inventor, if any	y:			A petitio	n has been file	d for this	unsign	ed inv	entor
Given Na	me (first and middle [if any])	)		Family Name or Surname						
Inventor's Signature				Date						
Residence: City		State		Country Citizenship						
Post Office Address										
Post Office Address										
City State					ZIP		Co	untry		,

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time—you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS SEND TO Assistant Commissioner for Patents, Washington, DC 20231.